arXiv:0809.2563v2 [math.DG] 16 Sep 2008

ON THE MEAN CURVATURE OF NASH ISOMETRIC EMBEDDINGS

G. PACELLI BESSA AND J. FABIO MONTENEGRO

Abstract. J. Nash proved in [7] that the geometry of any Riemannian manifold M imposes no restrictions to be embedded isometrically into a (fixed) ball $B_{\mathbb{R}^N}(1)$ of the Euclidean space $\mathbb{R}^N$. However, the geometry of M appears, to some extent, imposing restrictions on the mean curvature vector of the embedding.

1. Introduction

In 1956, John Nash in a celebrated article [7], proved that any complete n-dimensional Riemannian manifold M can be isometrically embedded in a ball $B_{\mathbb{R}^N}(1)$ of the Euclidean space $\mathbb{R}^N$ of radius 1 with $N = n(n+1)(3n+11)/2$. Although the geometry of M does not impose restrictions on the existence of an isometric embedding $\varphi : M \hookrightarrow B_{\mathbb{R}^N}(1)$, it does impose restrictions on the mean curvature vector of φ. That is expressed in the following theorem.

Theorem 1.1. *Let $\varphi : M \hookrightarrow B_{\mathbb{R}^N}(R) \subset \mathbb{R}^N$ be an isometric embedding of a complete n-dimensional Riemannian manifold into the Euclidean space $\mathbb{R}^N$. Then*

$$\sup_M \|\vec{H}\| \geq n/R - \lambda^*(M)\cdot R/2$$

Where $\lambda^*(M)$ is the fundamental tone of M given by

$$\lambda^*(M) = \inf\left\{\frac{\int_M |\nabla f|^2}{\int_M f^2},\, f \in H_0^1(M)\setminus\{0\}\right\},$$

$\vec{H}= \mathrm{tr}\alpha$ *is the mean curvature vector and α is the second fundamental form of φ.*

This result or rather its proof was already given in the literature, for instance in [2], [3], [4], [6] but these restrictions on the mean curvature vector of Nash embeddings was never observed.

2. Proof of Theorem 1.1

Theorem 1.1 is an application of the following variation of Barta's Theorem [1].

Theorem 2.1. *Let M be a complete Riemannian manifold. Let f be a positive smooth function on M. Then $\lambda^*(M) \geq \inf_M(-\triangle f/f)$.*

The proof of Theorem 1.1 is as follows. Let $\varphi : M \hookrightarrow B_{\mathbb{R}^N}(1) \subset \mathbb{R}^N$ be an isometric immersion. Let $g : \mathbb{R}^N \to \mathbb{R}$ defined by $g = (R^2 - \rho^2)/2$, where ρ is the distance function in $\mathbb{R}^N$ to the origin. Define $f = g\circ\varphi : M \to \mathbb{R}$. Observe that $f > 0$ thus by Barta's Theorem we have that $\lambda^*(M) \geq \inf_M(-\triangle f/f)$. To compute $\triangle f$ we use the following formula proved by Jorge-Koutrofiotis in [5].

$$\triangle f(y) = \sum_{i=1}^{n} \mathrm{Hess}\, g(\varphi(y)(e_i,e_i) + \langle \mathrm{grad}\, g, \vec{H}\rangle(\varphi(y)) \tag{2.1}$$

Date: September 16, 2008.

where $\{e_i\}$ is an orthonormal basis of T_yM identified with $\{d\varphi(y)e_i\}$.

Choose a basis $\{e_i\}$ as follows. Let $\{\operatorname{grad}\rho, \partial/\partial\theta_1, \dots \partial/\partial_{N-1}\}$ be a polar basis for $T_{\varphi(y)}\mathbb{R}^N$ and let $e_1 = \langle e_1, \operatorname{grad}\rho\rangle \operatorname{grad}\rho + \langle e_1, \operatorname{grad}\rho^{\perp}\rangle \operatorname{grad}\rho^{\perp}$ and $e_2, \dots, e_n \in \{\partial/\partial\theta_1, \dots \partial/\partial_{N-1}\}$. With this choice we obtain

$$\begin{aligned}
\triangle f &= \sum_{i=1}^{n} \operatorname{Hess} g(e_i, e_i) + \langle \operatorname{grad} g, \vec{H}\rangle \\
&= g''\langle e_1, \operatorname{grad}\rho\rangle^2 + \frac{g'}{\rho}\langle e_1, \operatorname{grad}\rho^{\perp}\rangle^2 + \frac{m-1}{\rho}g' + g'\langle \operatorname{grad}\rho, \vec{H}\rangle \\
&= (g'' - \frac{g'}{\rho})\langle e_1, \operatorname{grad}\rho\rangle^2 + \frac{m}{\rho}g' + g'\langle \operatorname{grad}\rho, \vec{H}\rangle \\
&= -n - \rho\langle \operatorname{grad}\rho, \vec{H}\rangle.
\end{aligned}$$

From this we have

$$-\frac{\triangle f}{f} = \frac{2n}{R^2-\rho^2} + \frac{2\rho}{R^2-\rho^2}\langle \operatorname{grad}\rho, \vec{H}\rangle \geq \frac{2n}{R^2} - \frac{2\sup_M \|\vec{H}\|}{R}. \tag{2.2}$$

We may assume that $\sup_M \|\vec{H}\| < \infty$, otherwise there is nothing to prove. We have then

$$\lambda^*(M) \geq \inf(-\triangle f/f) \geq \frac{2n}{R^2} - \frac{2\sup_M \|\vec{H}\|}{R}$$

Therefore

$$\sup_M \|\vec{H}\| \geq n/R - \lambda^*(M) \cdot R/2. \tag{2.3}$$

If we consider the mean curvature function H defined by $\vec{H} = n \cdot H \cdot \vec{\eta}$, where $\vec{\eta}$ is unit vector normal to the submanifold, the inequality (2.3) becomes $\sup_M |H| \geq 1/R - \lambda^*(M) \cdot R/2n$.

Remark 2.2. The hyperbolic n-space $\mathbb{H}^n(-1)$ has $\lambda^*(\mathbb{H}^n(-1)) = (n-1)^2/4$. Thus it can not be embedded (immersed) minimally into a ball $B_{\mathbb{R}^{N(n)}}(1)$ for $n = 2, \dots, 9$.

Acknowledgments: We would like to thank Paolo Piccione for the discussions we had with respect to this note when he visited us in Fortaleza.

REFERENCES

[1] J. Barta, *Sur la vibration fundamental d'une membrane.* C. R. Acad. Sci. **204**, (1937), 472–473.

[2] G. P. Bessa and J. Fabio Montenegro, *Eigenvalue estimates for submanifolds with locally bounded mean curvature.* Ann. Global Anal. and Geom. 24, 2003, 279-290.

[3] G. P. Bessa, and J. Fabio Montenegro, *An Extension of Barta's Theorem and Geometric Applications.* Ann. Global Anal. Geom. **31** (2007), no. 4, 345–362.

[4] G. P. Bessa, and M. Silvana Costa, *Eigenvalue Estimates for submanifolds with locally bounded mean curvature in $N \times \mathbb{R}$.*

[5] L. P. Jorge, and D. Koutroufiotis, *An estimative for the curvature of bounded submanifolds*, Amer. J. of Math. **103**, (1980), 711–725.

[6] L. P. Jorge and F. Xavier, *An inequality between the exterior diameter and the mean curvature of bounded immersions.* Math. Z. **178**, (1981) 77–82.

[7] J. Nash, *The imbedding problem for Riemannian manifolds.* Ann. of Math. (2) **63**,(1956) 20–63.